

Provided for non-commercial research and education use.

Not for reproduction, distribution or commercial use.



This article was published in an CASRP journal. The attached copy is furnished to the author for non-commercial research and education use, including for instruction at the authors institution, sharing with colleagues and providing to institution administration.

Other uses, including reproduction and distribution, or selling or licensing copied, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding CASRP's archiving and manuscript policies encouraged to visit:

<http://www.casrp.co.uk/journals>

© 2016 CASRP Publishing Company Ltd. UK.



Available online at www.casrp.co.uk/journals

CASRP PUBLISHER

**International Journal of Advanced Biological
and Biomedical Research 4(2) (2016) 129–135**

doi: 10.18869/IJABBR.2016.129



Original Article

Open Access

Effect of salinity on some physiological and biochemical traits in grape (*Vitisvinifera*)

Azar Seydi^{1,*}, Elham Faryabi², Farzad Kianersi³

¹M.Sc of Agricultural and Natural Resources Research and Education Center, Ilam, Iran.

²Department of Agriculture, Payame Noor University of Iran.

³Ph.D. Student of Plant Breeding Bu-Ali Sina, Hamedan, Iran.

Abstract

The present study was laid out in order to evaluate the effect of salinity on some biochemical characteristics of grapevine cv. Yaghouti. The experiment was done at Agricultural and Natural Resources Research Center, Ilam, Iran at 2014. The experiment was a completely block design (CRD) with four replications. Saline factor was NaCl salinity in seven levels (Zero, 50, 100, 150, 200 and 250 mM). Grape scions were planted in beds containing equal proportions of sand, perlite and vermiculite in 20-L vases. The vases were nourished by half Hoagland nutrient solution for first three weeks after plantation. In order to create the desired salinity levels, another half of the Hoagland nutrient solution was salinized with 0, 50, 100, 150, 200 and 250 mM NaCl. During growth stages, measurement was done on physiological and biochemical parameters such as soluble sugar, proline, chlorophyll, catalase, peroxidase and ascorbate peroxidase enzymes. The results showed that maximum chlorophyll a and b were obtained for control treatment and decreased gradually with increasing of saline level. The proline and soluble content as osmolites increased gradually by increasing of saline levels. However, CAT, APX and POX activity increased by salinity levels until 150mM, 150mM and 250mM saline treatment respectively. The results of the present study revealed that salt stress decreased some biochemical traits and decreased some damages by increasing of osmolites content and antioxidant enzymes activity.

© 2016 Published by CASRP publishing company Ltd. UK. Selection and/or peer-review under responsibility of Center of Advanced Scientific Research and Publications Ltd. UK.

Keywords: Enzyme, Chlorophyll, Proline.

* Corresponding author: M.Sc of Agricultural and Natural Resources Research and Education Center, Ilam, Iran.

© 2016 The Authors. This is an open access article under the terms of the Creative Commons Attribution-Non Commercial- No Derives License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Received 19 March 2016

Accepted 18 April 2016

Available online 25 April 2016

iThenticate screening 22 March 2016

English editing 15 April 2016

Quality control 21 April 2016

1. Introduction

Grapevines (*Vitisvinifera* L.) are one of the oldest and most important perennial crops in the world. An estimated 5000 to 6000 varieties of the *Vitisvinifera* species exist around the world, although only a relatively small number of these are used commercially (Dettweiler and Eibach, 2003). So far identify 30555 grapevine cultivars and many species on the world then naming and evaluated. Iran is one of the creation and distribution centers of grapes in the world that had high genetic diversity (Dettweiler and Eibach, 2003). Among the Iranian grapevine cultivars, only good quality cultivars for fresh grapes, raisins and industrial processing seem to have attracted farmers and others less attention.

Salinity has considerable adverse impacts on productivity of agricultural plants. Soil salinity, resulting from natural processes or from crop irrigation with saline water, occurs in many arid and semi-arid regions of the world (Koca et al., 2007). Soil salinity affects growth and yield in grapevine by osmotic and specific ion toxicities (Shani and Al, 2005). One of the most common stress responses in high plants is over production of some various compatible organic solutes (Serraj and Sinclair, 2002). These solutes are most commonly carbohydrates, like sugars, amino acids and proteins that acts as osmolytes. Proline and soluble sugar are two of them that known to occur widely in higher plants and normally accumulates in large quantities in response to environmental stresses (Kishor et al., 2005). It has also been reported that not only species but also varieties of a species show differences in salt tolerance (Schwarz, 1995). In plant cells, antioxidant enzymes, such as superoxide dismutase (SOD), peroxidase (POX), and catalase (CAT), are considered to form a defensive team, whose combined purpose is to protect cells from oxidative damage (Blokina et al., 2003). Moreover, Goel and Sheoran (2003) told that malondialdehyde (MDA) is considered sensitive marker commonly used for assessing membrane lipid peroxidation. CAT reduces H₂O₂ to water and dioxygen, and APX reduces H₂O₂ water and generates MDA (Noctor and Foyer, 1998). In a study, three European vinifera cultivars and four Argentinean cultivars were analyzed for their salt tolerance by in vitro culture and it was found that Argentinean cultivars showed better performance in growing under salinity, especially in the highest salt concentration (Cavagnaro et al., 2006). However, this study was conducted to study on some biochemical characterize of grapevine under salt stress.

2. Materials and methods

This experiment was laid out in order to evaluate the effect of salinity on some physiological and biochemical of grape (*Vitisvinifera*) c.vYaghoutiin Agricultural and Natural Resources Research Center, Ilam, Iran at 2014. The experiment was a completely block design (CRD) with four replications at greenhouse. Saline factor was NaCl salinity in seven levels (Zero, 50, 100, 150, 200 and 250 mM). Grape scions were planted in beds containing equal proportions of sand, perlite and vermiculite in 20-L vases. The vases were nourished by half Hoagland nutrient solution for first three weeks after plantation. In order to create the desired salinity levels, another half of the Hoagland nutrient solution was salinized with 0, 50, 100, 150, 200 and 250 mMNaCl.

The vases were irrigated by the prepared solutions after the establishment of scions. During growth stages, measurement was done on physiological and biochemical parameters such as soluble sugar, proline, chlorophyll, catalase, peroxidase and ascorbat peroxidase enzymes. Soluble protein was estimated using the bovine serum albumin as standard. Extraction of protein from the sample was carried out using tris HCl buffer (pH 7.8) and the supernatant was used for the estimation and the absorbance was measured at 660 nm (Sadasavim et al., 1966).

To measure proline content, cotyledon (500 mg) were ground in mortal and pistil, homogenized in 10 ml 3% sulfosalicylic acid, and then centrifuged at 5 000 rpm for 5 min. The supernatant (200 µl) was mixed with 200 µl glacial acetic acid and 200 µl acidic ninhydrin, and then mixed well. The mixture was incubated at 100°C for 60 min. The reaction was terminated by placing the mixture on ice, and then extracting the sample with 1.0 ml toluene. The absorbance of the mixture was measured at 520 nm. To prepare the standard curve, an L-proline stock solution (100 µg/ml) was diluted to 2, 4, 6, 8, 10, 12 µg/ml (Bates, 1973).

Chlorophyll a and b was measured using Arnonmethod. In this method, as little as a 0.5 gram of wet vegetative matter was mashed in liquid nitrogen, in a porcelain mortar. As much as 20mL of 80% acetone was added to the sample, and then the mixture was put into centrifuge device with 6000 rpm speed for 10 minutes. Supernatant was transferred into a glass balloon. Some of the samples in the balloon were read in spectrophotometer for chlorophyll a at 663nm; for chlorophyll b at 645nm; and for Carotenoids at 470nm. Finally,

the following formulas were used to calculate chlorophyll a and b and carotenoids contents in mg/g of fresh weight of the sample:

$$\text{Chlorophyll a} = (19.3 * A_{663} - 0.86 * A_{645}) V/100W$$

$$\text{Chlorophyll b} = (19.3 * A_{645} - 3.6 * A_{663}) V/100W$$

For the catalase (CAT) assay, soluble proteins were extracted by homogenizing 1 g (fresh weight) powdered sample in 3 ml of 100 mM mono sodium phosphate buffer (pH 7.5) containing 0.1% Triton X-100. Other materials for determined of CAT activity was 54 mM mono sodium phosphate buffer (pH 6.8) and 450 mM H₂O₂. CAT activity was measured following the change in the absorbance of the reaction mixture at 240 nm (Philippe et al., 2007). The assay was detected in a 3 ml reaction mixture containing 2.85 mL 54 mM mono sodium phosphate buffer (pH 6.8), 100 µL H₂O₂ (450 mM) and 50 µL of crude extract.

POX activity was measured by Sakharov and Aridilla (1999) method with slight modifications. A 3 mL mixture consisted of 100 µL of based solution guaiacol, 100 µL H₂O₂ (1.2 M) and 200 µL enzyme extract and the changes of absorbance at 470nm were measured using a UV/vis spectrophotometer. One unit of POX activity was expressed as 1.0 change in absorbance per minute. Total APX as Nakano and Asada (1981) method with slight modifications in a 1-ml total volume. The reaction mixtures included 25 µL ascorbic acid (0.03%), 25 µL EDTA (0.1%), 50 µL H₂O₂ (0.015%), 800 µL 312.5 mM sodium phosphate buffer (pH 7.0) and 100 µL protein extract. The changes of absorbance were read at 290 nm using a UV/vis spectrophotometer for 3 minutes.

The statistical analysis was conducted using JMP 5.0.1.2 (Statistical analyses system Institute incorporated, 2002). Statistical significance was declared at P≤0.05 and P≤0.01. Treatment effects from the two runs of experiments followed a similar trend, and thus the data from the two independent runs were combined in the analysis.

3. Results and discussion

The results showed that some physiological and biochemical traits of grape Yaghuti cultivar was changed by increasing of saline treatment. The results showed that effect of saline treatment on all traits was significant. However, mean comparison results showed that maximum chlorophyll-a was recorded for control treatment (8.1 mg/ml). Minimum chlorophyll-b was recorded for 250 mM saline treatment (2.72 mg/ml) (fig1).




Fig. 1. Effect of salinity on chlorophyll-a in *vitisvinifera* leaves (Columns with same letters have not significant differences).

The results showed that maximum chlorophyll-b was recorded for control treatment (0.84 mg/ml). Minimum chlorophyll-b was recorded for 250 mM saline treatment (0.13 mg/ml) (fig2).




Fig. 2. Effect of salinity on chlorophyll –b in *vitisvinifera* leaves (Columns with same letters have not significant differences).

The results of the present study showed that maximum soluble sugar content was recorded for 250 mM saline treatment (366 mg/g). Minimum soluble sugar content was recorded for control treatment (146 mg/g) (fig3).




Fig. 3. Effect of salinity on soluble sugar content in *vitisvinifera* leaves (Columns with same letters have not significant differences).

Proline content as an osmolit increased gradually by increasing of saline levels until 250mM. However, maximum proline content recorded for 250 mM saline treatment (270 mg/g). Minimum proline content was recorded for control treatment (86 mg/g) (fig4).




Fig. 4. Effect of salinity on proline content in *vitisvinifera* leaves (Columns with same letters have not significant differences).

Activity of CAT enzyme decreased by salinity levels until 150mM saline treatment then decreased. However, maximum CAT recorded for 150 mM saline treatment (2630 nMol.mn⁻¹.g⁻¹ FW). Minimum CAT activity was recorded for control treatment (1860 nMol.mn⁻¹.g⁻¹ FW) (fig5).




Fig. 5. Effect of salinity on CAT activity in *vitisvinifera* leaves (Columns with same letters have not significant differences).

POX activity increased by salinity levels until 250 mM saline treatment. However, maximum POX recorded for 250mM saline treatment (121 nMol.mn⁻¹.g⁻¹ FW). Minimum POX activity was recorded for control treatment (61 nMol.mn⁻¹.g⁻¹ FW) (fig6).




Fig. 6. Effect of salinity on POX activity in *vitisvinifera* leaves (Columns with same letters have not significant differences).

APX activity increased by salinity levels until 150mM saline treatment then decreased. Maximum APX recorded for 150 mM saline treatment (69 nMol.mn⁻¹.g⁻¹ FW). Minimum APX activity was recorded for control treatment (31 nMol.mn⁻¹.g⁻¹ FW) (fig7).




Fig. 7. Effect of salinity on APX activity in *vitisvinifera* leaves (Columns with same letters have not significant differences).

The results of the present study showed that saline treatment had the significant effect on physiological and biochemical parameters such as soluble sugar, proline, chlorophyll, catalase, peroxidase and ascorbat peroxidase enzymes. Chlorophyll a and b were decreased by increasing of saline treatment (fig 1 and 2). One of the remarkable effects of saline treatment is decline of chlorophyll a and b. Increasing salinity level leads to decreased chlorophyll biosynthesis through increased salt. Increasing of saline treatment, lead to increasing of some osmolites such as proline and soluble sugar content (fig 3 and 4). Under high salt stress plant cells decrease their osmotic potential by accumulation of some solutes such as proline and soluble sugars (Youssef and Al-Fredan, 2008). Proline plays an adaptive role in mediating osmotic adjustment and protecting the sub-cellular structures in stressed plants (Voetberg and Sharp, 1991). Proline is more efficiency to protection against stress comparison with other conventional osmolit compatible such as sugars and sugar alcohols (Verma, 2003). Some enzymes such as CAT, POX and APX had the anti-oxidative roles in front of oxidative stresses. In the present study, results showed that with increasing of saline treatment until 150 mM CAT and APX activities increased than decreased. However POX activity was increased by increasing of saline treatment significantly. Salt stresses could cause oxidative damage. Therefore, plant cells need different mechanisms, which enable the detoxification of excess reactive oxygen species (ROS) and keep the balance of the formation and removal ROS. The antioxidant enzymes remove some dangerous reactive oxygen species produced by oxidative stresses. The imbalance between the production of reactive species and antioxidants is called oxidative stress; this condition is known as a cause of damage in biomolecules (DNA, lipids and proteins) and is involved in the generation or aggravation of more than a hundred pathological conditions (Berger, 2005). In this study, the changing of CAT activity was related with the level of salinity. There were significant changes in CAT, POX and APX activities with increasing of salinity levels as oxidative stress. In final the results of the present study reviled that salt stress decreased some biochemical traits and decreased some damages by increasing of proline and sugar content and antioxidant enzymes such as CAT, POX and APX.

References

- Bates, I.S., Waldern, R.P., Teare, I.D., 1973. Rapid determination of free proline for water stress studies. *Plant. Soil.*, 39, 905-207.
- BERGER, M.M., 2005. Can oxidative damage be treated nutritionally. *Clin. Nutr.*, 24, 172-183.
- Blokhina, O., Virolainen, E., Fagerstedt, K.V., 2003. Antioxidants, oxidative damage and oxygen deprivation stress: a review. *Ann. Botan.*, 91, 179–194.
- Cavagnaro, J.B., Ponce, M.T., Guzman, J., Cirrincione, M.A., 2006. Argentinean cultivars of *Vitis vinifera* grow better than European ones when cultured in vitro under salinity. *Biocell.*, 30(1), 1-7.
- Dettweiler, E., Eibach, R., 2003. The two *Vitis* databases as tools for germplasm management *Vitis international variety catalogue and European Vitis database*. *Acta. Horticulture.*, 603, 505-509.
- Kishor, P.B.K., Sangam, S., Amrutha, R.N., Laxmi, P.S., Naidu, K.R., Rao, K.S., 2005. Regulation of proline biosynthesis, degradation, uptake and transport in higher plants: its implications in plant growth and abiotic stress tolerance. *Curr. Sci.*, 88, 424-38.
- Koca, M., Bor, M., Ozdemir, F., Turkan, I., 2007. The effect of salt stress on lipid peroxidation, antioxidative enzymes and proline content of sesame cultivars. *Environ. Exp. Bot.*, 60, 344–351.
- Nakano, Y., Asada, K., 1981. Hydrogen peroxide scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant. Cell. Physiol.*, 22, 867–880.
- Noctor, G., Foyer, C.H., 1998. Ascorbate and glutathione: keeping active oxygen under control. *Ann. Rev. Plant. Physiol. Plant. Molecul. Biol.*, 49, 249–279.
- Philippe, M., Koraljka, R.K., Karlheinz, B., 2007. A simple method to measure effective catalase activities: optimization, validation, and application in green coffee. *Analytic. Biochem.*, 360, 207–215.
- Sadasivam, N., Manickam, A., 1996. *Biochemical Methods for Agricultural Sciences*, New Age International (P) Ltd., New Delhi.
- Schwarz, M., 1995. Soilless culture management. *Adv. Ser. Agr. Sci.*, 24, 197.
- Serraj, R., Sinclair, T.R., 2002. Osmolyte accumulation: can it really help increase crop yield under drought conditions? *Plant. Cell. Environ.*, 25, 333-341.
- Shani, U., Ben-Gal, Al., 2005. Long-term response of grapevines to salinity: Osmotic effects and ion toxicity. *Am. J. Enol. Vitic.*, 56, 148-154.

- Voetberg, G.S., Sharp, R.E., 1991. Growth of the maize primary root in low water potentials. III. Roles of increased praline depositions in osmotic adjustment. *Plant. Physiol.*, 96, 125-30.
- Youssef, A.M., Al-Fredan, M.A., 2008. Community composition of major vegetations in the coastal area of Al-Uqair, Saudi Arabia in response to ecological variations. *J. Biol. Sci.*, 8, 713-721.

How to cite this article: Seydi, A., Faryabi, E., Kianersi, F., 2016. Effect of salinity on some physiological and biochemical traits in grape (*Vitisvinifera*). *International journal of Advanced Biological and Biomedical Research*, 4(2), 129-135.

Submit your next manuscript to CASRP Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.casrp.co.uk/journals

